VETERINARY MEDICINE

Research of Endocrinology

Prepared Under the Supervision of

Dr. Sherif Elmosalamy

Prepared By

Manal Gomaa Rabea	592500739
Farah Ahmed Hussien	592500433
Shahd Mohamed Hammad	592500458

Academic Research Paper

Endocrinology is the branch of medicine concerned with hormones, the glands that produce them, and disorders caused by hormone excess, deficiency, or abnormal regulation. In veterinary medicine, it focuses on how hormones maintain homeostasis and coordinate metabolism, growth, reproduction, stress responses, and fluid and electrolyte balance in animals.

balance in animals.

faster electrical and chemical signals through neurons. Both systems work together to maintain the internal stability of the body.

Clinical importance in dogs and cats

Endocrine disorders are common in companion-animal practice. Important examples include diabetes mellitus, hyperthyroidism in cats, hypothyroidism in dogs, hyperadrenocorticism (Cushing's syndrome), and hypoadrenocorticism (Addison's disease). These conditions may affect several body systems and can produce signs that are initially subtle or nonspecific.

A sound understanding of normal endocrine physiology helps veterinarians recognize disease patterns, select appropriate laboratory tests, interpret results in context, and choose rational treatment. Advances in laboratory testing, diagnostic imaging, and medicines have improved the long-term management of many endocrine disorders.

1.1 Organization of the Endocrine System

The endocrine system includes distinct glands and hormone-producing cells within other organs. The major glands are the hypothalamus, pituitary, thyroid, parathyroids, adrenal glands, endocrine pancreas, and gonads. The hypothalamus and pituitary form a central control unit: hypothalamic signals regulate pituitary secretion, and pituitary hormones stimulate peripheral endocrine glands.

Endocrine activity also occurs in the gastrointestinal tract, heart, kidneys, adipose tissue, and placenta. These organs release signals that contribute to digestion, blood pressure regulation, red-cell production, energy balance, and reproduction.

1.2 Major Functions

• Metabolism: thyroid hormones, insulin, glucagon, and cortisol.

• Water and electrolyte balance: aldosterone and antidiuretic hormone (ADH).

• Stress response: cortisol and catecholamines.

• Growth and development: growth hormone and thyroid hormones.

• Reproduction: gonadotropins and sex steroids.

2. Major Endocrine Glands and Their Locations

The major endocrine glands are distributed throughout the body. Their hormones act on specific target tissues and coordinate several physiological functions.




Figure 1. Major endocrine glands in the cat, including the hypothalamus, pituitary, thyroid, parathyroids, adrenals, pancreas, and gonads.

Gland	Location / Main role
Hypothalamus	Ventral region of the brain; integrates neural and endocrine signals.
Pituitary gland	At the base of the brain; regulates several peripheral endocrine glands.
Thyroid gland	Two lobes beside the trachea; produces T3, T4, and calcitonin.
Parathyroid glands	Small glands associated with the thyroid; secrete PTH.
Adrenal glands	Paired organs cranial to the kidneys; produce steroid hormones and catecholamines.
Endocrine pancreas	Islets of Langerhans; secrete insulin, glucagon, and other hormones.
Gonads	Ovaries and testes; produce sex steroids and support reproduction.

3. Hormones: Types and Functions

Hormones can be classified according to their chemical structure. Their solubility influences how they circulate in blood and how they interact with target cells.

3.1 Main Chemical Classes

• **Peptide and protein hormones:** Examples include insulin, growth hormone, ACTH, TSH, FSH, LH, and PTH. They are water-soluble and usually bind to receptors on the cell membrane.

• **Steroid hormones:** Cortisol, aldosterone, estrogen, progesterone, and testosterone are derived from cholesterol. They are lipid-soluble, often circulate bound to carrier proteins, and act through intracellular receptors.

• **Amino-acid-derived hormones:** Catecholamines (epinephrine and norepinephrine) and thyroid hormones (T3 and T4) belong to this group. Catecholamines act mainly through membrane receptors, while thyroid hormones act mainly through nuclear receptors.

3.2 Mechanisms of Hormonal Action

Membrane-receptor pathway: Peptide hormones and catecholamines bind to receptors on the cell surface. This activates intracellular signaling pathways and second messengers such as cAMP, IP3, DAG, calcium, or kinase cascades. These pathways can rapidly change enzyme activity or ion-channel function.

Nuclear-receptor pathway: Steroid and thyroid hormones enter cells and bind intracellular receptors. The hormone–receptor complex influences gene transcription, producing effects that may take longer to develop but can last longer.

The number and sensitivity of receptors can change in response to hormone concentrations. This regulation helps tissues adjust their responsiveness and is important when interpreting endocrine tests and treatment responses.

3.3 Hormone–Target Cell Specificity

A hormone does not affect every cell equally. Its action depends on the presence of the appropriate receptor and the signaling machinery within the target cell. Differences in receptor distribution help explain why one hormone can produce different effects in different organs.

Clinical note: hormone test results should be interpreted alongside the animal’s clinical signs, concurrent illness, medication history, and the specific assay used.

4. Hypothalamus and Pituitary Gland

The hypothalamus links the nervous and endocrine systems. It receives information about the animal's internal state, blood chemistry, and biological rhythms, then regulates pituitary activity through releasing and inhibiting hormones.

4.1 Hypothalamic Hormones

• TRH stimulates thyroid-stimulating hormone (TSH) release.

• CRH stimulates adrenocorticotropic hormone (ACTH) release.

• GnRH stimulates follicle-stimulating hormone (FSH) and luteinizing hormone (LH).

• GHRH stimulates growth hormone (GH), while somatostatin inhibits GH release.

• Dopamine inhibits prolactin secretion.

4.2 Anterior Pituitary

The anterior pituitary produces several hormones that regulate growth, metabolism, reproduction, and peripheral endocrine glands:

• **ACTH:** stimulates the adrenal cortex.

• **TSH:** stimulates thyroid hormone production.

• **GH:** promotes growth and influences metabolism, partly through insulin-like growth factor 1 (IGF-1).

• **FSH and LH:** regulate gonadal function and reproduction.

• **Prolactin:** supports lactation.

4.3 Posterior Pituitary

The posterior pituitary stores and releases hormones synthesized in the hypothalamus. ADH (vasopressin) supports water conservation by the kidneys, while oxytocin contributes to uterine contraction and milk ejection.

Disorders involving the hypothalamic–pituitary axis may cause multiple hormone deficiencies or excesses. Examples include pituitary-dependent hyperadrenocorticism, pituitary dwarfism, and central diabetes insipidus.




Figure 3. Location and main anatomical parts of the pituitary gland.

5. Thyroid and Parathyroid Glands

5.1 Thyroid Gland

The thyroid gland produces thyroxine (T4) and triiodothyronine (T3), iodine-containing hormones that influence basal metabolic rate, protein synthesis, growth, and the function of many organ systems. Parafollicular C-cells produce calcitonin, which contributes to lowering blood calcium.

Thyroid hormones affect nearly every tissue. Therefore, both deficiency and excess can produce signs involving several body systems. In dogs, hypothyroidism is a common endocrinopathy. In older cats, hyperthyroidism is especially prevalent and is commonly associated with benign adenomatous hyperplasia.

5.2 Hypothyroidism in Dogs

Canine hypothyroidism is commonly associated with lymphocytic thyroiditis or idiopathic thyroid atrophy. Clinical signs may include lethargy, weight gain, cold intolerance, hair loss, skin changes, and reduced activity. Diagnosis is based on clinical assessment and thyroid testing, commonly total or free T4 with TSH interpreted in context. Treatment generally involves lifelong levothyroxine supplementation.

5.3 Hyperthyroidism in Cats

Feline hyperthyroidism is common in older cats. Typical signs include weight loss despite increased appetite, increased activity, rapid heart rate, vomiting, and increased thirst and urination. Diagnosis is usually supported by an elevated serum total T4; free T4 may help in selected borderline cases. Treatment options include antithyroid medication, thyroidectomy, radioactive iodine, or an iodine-restricted diet.

5.4 Parathyroid Glands

The parathyroid glands secrete parathyroid hormone (PTH) in response to low blood calcium. PTH raises serum calcium by increasing bone resorption, enhancing renal calcium reabsorption, and stimulating vitamin D activation, which increases intestinal calcium absorption. It also promotes phosphate excretion.

Thyroid test interpretation requires care because concurrent illness can lower total T4 without primary thyroid disease. This is known as non-thyroidal illness or euthyroid sick syndrome.

6. Adrenal Glands

The adrenal glands are paired organs located cranial to the kidneys. Each gland has an outer cortex and an inner medulla. The cortex produces steroid hormones, while the medulla produces catecholamines.

6.1 Adrenal Cortex

• **Zona glomerulosa:** produces mineralocorticoids, mainly aldosterone, which regulate sodium, potassium, and water balance.

• **Zona fasciculata:** produces glucocorticoids, mainly cortisol, which influence metabolism, stress responses, and immune function.

• **Zona reticularis:** produces smaller quantities of sex steroids, including androgens and estrogens.

6.2 Adrenal Medulla

The adrenal medulla secretes epinephrine and norepinephrine in response to sympathetic nervous system stimulation. These catecholamines contribute to the rapid “fight-or-flight” response, including changes in heart rate, blood flow, and energy availability.

6.3 Hyperadrenocorticism (Cushing’s Syndrome)

Hyperadrenocorticism results from excessive glucocorticoid exposure. In dogs, many cases are pituitary-dependent due to excess ACTH, while a smaller proportion are caused by an adrenal tumor. Common signs include increased thirst and urination, increased appetite, abdominal enlargement, panting, muscle weakness, hair loss, and thin skin. Diagnostic tests may include the low-dose dexamethasone suppression test, urine cortisol:creatinine ratio, endogenous ACTH, and imaging. Trilostane is commonly used for medical management.

6.4 Hypoadrenocorticism (Addison’s Disease)

Hypoadrenocorticism is caused by inadequate adrenal hormone production. Primary adrenal failure can result in deficiency of both cortisol and aldosterone. Signs may include lethargy, poor appetite, weight loss, and intermittent gastrointestinal problems. Severe cases may develop an Addisonian crisis with dehydration and electrolyte abnormalities. Diagnosis is confirmed using an ACTH stimulation test. Long-term treatment involves glucocorticoid and, when needed, mineralocorticoid replacement.

7. Endocrine Pancreas and Reproductive Endocrinology

7.1 Endocrine Pancreas

The endocrine pancreas contains clusters of cells called the islets of Langerhans. Different cell types produce hormones that regulate blood glucose and digestive activity:

• **Beta (β) cells:** insulin, which lowers blood glucose by promoting uptake and storage of nutrients.

• **Alpha (α) cells:** glucagon, which raises blood glucose by stimulating hepatic glycogen breakdown and glucose production.

• **Delta (δ) cells:** somatostatin, which modulates secretion of other hormones.

• **PP cells:** pancreatic polypeptide, which contributes to regulation of pancreatic and gastrointestinal functions.

7.2 Diabetes Mellitus

Diabetes mellitus is a frequent endocrine disease in dogs and cats. Dogs commonly develop an insulin-deficient, type 1-like condition associated with loss of beta-cell function. Cats more often develop a type 2-like condition involving insulin resistance and relative insulin deficiency, although many eventually require insulin therapy.

Typical clinical signs include increased thirst, increased urination, increased appetite, and weight loss. Diagnosis is based on persistent hyperglycemia together with glucosuria; fructosamine can help assess whether hyperglycemia is sustained. Treatment usually includes insulin, dietary management, and regular monitoring for complications such as ketoacidosis, hypoglycemia, and urinary tract infection.

7.3 Reproductive Endocrinology

The hypothalamic–pituitary–gonadal axis regulates reproductive function. GnRH stimulates pituitary release of FSH and LH. In females, FSH supports follicular development and estrogen production, while LH triggers ovulation and supports the corpus luteum and progesterone production. In males, FSH supports sperm production and LH stimulates testosterone secretion by Leydig cells.

Sex steroids provide feedback to the hypothalamus and pituitary. Reproductive endocrine disorders may include persistent estrus, hormone-producing testicular tumors, and hormonal changes associated with reproductive disease or gonadectomy.

8. Feedback Mechanisms and Environmental Influences

8.1 Negative Feedback

Most endocrine axes are regulated by negative feedback. When the concentration of a peripheral hormone rises, it inhibits the hypothalamic and/or pituitary signals that stimulate its production. For example, increased cortisol reduces CRH and ACTH drive. This mechanism helps keep hormone concentrations within an appropriate physiological range.

8.2 Positive Feedback

Positive feedback is less common. An example occurs before ovulation, when rising estrogen temporarily stimulates rather than inhibits hypothalamic and pituitary activity, contributing to the LH surge. Understanding feedback is important when interpreting dynamic tests such as ACTH stimulation and dexamethasone suppression tests.

8.3 Environmental and Physiological Influences

• **Stress and illness:** may increase cortisol and reduce thyroid hormone measurements.

• **Nutrition and body condition:** influence insulin sensitivity, growth-related hormones, and reproductive function.

• **Photoperiod and season:** can affect reproductive cycles and melatonin secretion in some species.

• **Age:** several endocrine diseases become more common in middle-aged and older animals.

• **Medications:** glucocorticoids, phenobarbital, sulfonamides, and other drugs may alter hormone concentrations or test interpretation.

Veterinarians should interpret laboratory findings in light of the animal’s current illness, medication history, age, nutritional status, and physiological condition. A laboratory result should not be interpreted in isolation.

9. Common Endocrine Diseases in Dogs and Cats

9.1 Canine Hypothyroidism

Canine hypothyroidism is usually associated with lymphocytic thyroiditis or idiopathic thyroid atrophy. Common findings include lethargy, weight gain, cold intolerance, dermatological changes, and hair loss. Diagnosis combines clinical findings with thyroid hormone testing. Levothyroxine is the usual long-term treatment, with follow-up testing used to assess response.

9.2 Feline Hyperthyroidism

Feline hyperthyroidism is most often caused by benign adenomatous thyroid changes. Affected cats may lose weight despite eating more, become unusually active, develop tachycardia, vomit, or drink and urinate more frequently. Serum total T4 is the usual first-line diagnostic test. Treatment choices include methimazole or carbimazole, thyroidectomy, radioactive iodine, and selected iodine-restricted diets.

9.3 Clinical Examination

A careful physical examination is essential because endocrine disorders may initially produce nonspecific signs. Body condition, skin and coat, heart rate, hydration, abdominal contour, appetite, water intake, and urination history can help guide further testing.



Figure 3. Veterinary examination of a dog. Clinical findings help guide the selection and interpretation of endocrine tests.

10. Additional Endocrine Disorders

10.1 Canine Hyperadrenocorticism

Cushing’s syndrome is associated with excessive cortisol exposure. Many canine cases are pituitary-dependent, while others arise from an adrenal tumor. Clinical signs can include increased thirst and urination, increased appetite, abdominal enlargement, panting, muscle weakness, hair loss, and fragile skin. Screening may involve urine cortisol:creatinine ratio or low-dose dexamethasone suppression. Additional tests and imaging help determine the underlying form. Trilostane is a commonly used medical treatment.

10.2 Canine Hypoadrenocorticism

Addison’s disease results from inadequate adrenal hormone production. Clinical signs may be vague and intermittent, including reduced energy, poor appetite, weight loss, and gastrointestinal upset. An acute crisis can be life-threatening and requires urgent veterinary care. Diagnosis is confirmed by ACTH stimulation testing. Long-term management generally requires glucocorticoid replacement and mineralocorticoid replacement when indicated.

10.3 Diabetes Mellitus

Diabetes mellitus commonly presents with increased thirst, urination, and appetite, often accompanied by weight loss. Persistent hyperglycemia and glucosuria support the diagnosis. Fructosamine may help confirm sustained hyperglycemia. Treatment is based on insulin therapy, appropriate dietary management, and ongoing monitoring for complications.

Clinical priority

Acute deterioration, collapse, severe dehydration, suspected ketoacidosis, or a possible Addisonian crisis requires prompt veteri edication administration, monito



Figure 4. Veterinary examination of a dog in a clinical setting.

11. Diagnosis, Treatment and Clinical Management

11.1 Diagnostic Approach

Diagnosis follows a structured sequence that combines history, physical examination, routine laboratory work, specific endocrine tests, and imaging when indicated.

• **Clinical suspicion:** assess age, breed, history, appetite, thirst, urination, body condition, skin changes, and physical findings.

• **Minimum database:** complete blood count, serum biochemistry, and urinalysis may reveal supportive abnormalities such as hyperglycemia, glucosuria, electrolyte changes, or increased alkaline phosphatase.

• **Specific endocrine testing:** total/free T4 and TSH for thyroid disorders; ACTH stimulation for Addison's disease and selected monitoring; low-dose dexamethasone suppression for Cushing's screening; glucose and fructosamine for diabetes.

• **Imaging:** ultrasonography of the thyroid or adrenal glands, thyroid scintigraphy, and CT or MRI in selected pituitary cases.

11.2 Treatment Principles

• **Hormone replacement:** levothyroxine for hypothyroidism, insulin for diabetes, and glucocorticoid/mineralocorticoid replacement for Addison's disease.

• **Reduction of excess hormone action:** trilostane for hyperadrenocorticism and methimazole or radioactive iodine for feline hyperthyroidism.

• **Surgery or ablative treatment:** thyroidectomy, adrenalectomy, or pituitary-directed treatment in selected cases.

• **Dietary management:** tailored nutritional plans may support management of diabetes and selected thyroid disorders.

11.3 Long-Term Monitoring and Owner Education

Many endocrine diseases require lifelong follow-up. Monitoring may include repeat clinical examinations, laboratory testing, assessment of treatment response, and adjustment of medication. Owners should understand the chronic nature of many endocrine disorders, how to administer prescribed medicines, and the importance of scheduled rechecks. They should also know the warning signs that require urgent veterinary attention.

12. Conclusion

The endocrine system is an integrated network that is essential for homeostasis, metabolism, growth, reproduction, and adaptation to stress. Endocrine disorders are common in dogs and cats and may affect multiple organ systems. A strong understanding of normal gland function, hormone action, and feedback regulation supports accurate clinical assessment and appropriate diagnostic testing.

The management of endocrine disease depends on identifying the underlying disorder, selecting evidence-based treatment, monitoring the patient over time, and educating caregivers. With appropriate veterinary care and follow-up, many animals with endocrine disorders can maintain a good quality of life for extended periods.

References

The following selected references support the physiology, clinical features, diagnosis, and management discussed in this paper.

1. Feldman EC, Nelson RW, Reusch C, Scott-Moncrieff JCR. *Canine and Feline Endocrinology*. 4th ed. Elsevier Saunders; 2015.

2. Mooney CT, Peterson ME, Shiel RE. *BSAVA Manual of Canine and Feline Endocrinology*. 5th ed. British Small Animal Veterinary Association; 2023.

3. Bugbee A, et al. 2023 AAHA Selected Endocrinopathies of Dogs and Cats Guidelines. *J Am Anim Hosp Assoc*. 2023;59(3):113–135. doi:10.5326/JAAHA-MS-7368.

4. Merck Veterinary Manual. Endocrine System. Merck & Co.; 2024. https://www.merckvetmanual.com/endocrine-system

5. Rijnberk A, Kooistra HS. *Clinical Endocrinology of Dogs and Cats: An Illustrated Text*. 2nd ed. Schlütersche; 2010.

6. Behrend EN, Kooistra HS, Nelson R, Reusch CE, Scott-Moncrieff JC. Diagnosis of spontaneous canine hyperadrenocorticism: 2012 ACVIM consensus statement. *J Vet Intern Med*. 2013;27(6):1292–1304.

7. Peterson ME. Hyperthyroidism in cats: What's causing this epidemic of thyroid disease and can we prevent it? *J Feline Med Surg*. 2012;14(11):804–818.

8. Reusch CE. Feline diabetes mellitus. In: Feldman EC, Nelson RW, Reusch C, Scott-Moncrieff JCR, eds. *Canine and Feline Endocrinology*. 4th ed. Elsevier; 2015.

9. Lathan P, Thompson AL. Management of hypoadrenocorticism (Addison's disease) in dogs. *Vet Med (Auckl)*. 2018;9:1–10.

10. Nelson RW. Canine hyperadrenocorticism. In: Ettinger SJ, Feldman EC, Côté E, eds. *Textbook of Veterinary Internal Medicine*. 8th ed. Elsevier; 2017.